

Relationship between subsurface structure and large-scale fissures in the northwestern region in Aso valley caused by the 2016 Kumamoto earthquake

*土井 一生¹、釜井 俊孝¹、後藤 聡²、東 良慶³、大倉 敬宏⁴、村尾 英彦⁵、美馬 健二⁶

*Issei Doi¹, Toshitaka Kamai¹, Satoshi Goto², Ryohei Azuma³, Takahiro Ohkura⁴, Hidehiko Murao⁵, Kenji Mima⁶

1. 京都大学防災研究所、2. 山梨大学大学院総合研究部、3. 大阪工業大学工学部、4. 京都大学大学院理学研究科、5. 村尾地研、6. 太田ジオリサーチ

1. Disaster Prevention Research Institute, 2. Integrated Graduate School of Medicine, Engineering, and Agricultural Sciences, Yamanashi University, 3. Faculty of Engineering, Osaka Institute of Technology, 4. Graduate School of Science, Kyoto University, 5. Murao Chiken, 6. Ohta Geo Research. Co., Ltd.

In accompanied with the 2016 Kumamoto earthquake, fissures with the length and the height a few hundred meters and two meters, respectively, emerged in Aso valley (inside the Aso caldera). In this area, the top layer consists of sediments in the caldera lake created by Aso-4 eruption with the thickness a few tens of meters. InSAR and seismic data show that the region with the size of 1-4 square meter moved 1-2 m northward horizontally during strong motion of the mainshock (Fujiwara et al., 2016; Doi et al., 2016). We investigated the mechanism of the movement of these regions by estimating subsurface structure beneath this area.

Two station spatial auto-correlation (2ST-SPAC) method (Hayashi and Craig, 2016) was applied to estimate subsurface structure using ambient noises. We succeeded to estimate the S-wave velocity structure to the depth of 130 m in and around the regions with fissures. In the regions where large scale fissures were developed, a layer with S-wave velocity less than 150 m/s lay from the surface to the depth of 60 m, followed by two layers with 250 m/s and 300 m/s at depths of 60-90 m and 90-130 m, respectively. This low-velocity layer was considered to represent soft sediments in the caldera lake due to Aso-4 eruption and consistent with the nearby boring profile. Two relatively higher layers might correspond to lava layers after Aso-4 eruption. Moreover, the S-wave velocity at the top surface to the depth of 5 m was so slow as 80 m/s. We continue to estimate the distribution of the soft sediments and lava structure beneath them, to elucidate how fissures were generated in this area.